

Goals and Benefits with HCV Treatment

This is a PDF version of the following document:

Module 4: [Evaluation and Preparation for Hepatitis C Treatment](#)

Lesson 1: [Goals and Benefits with HCV Treatment](#)

You can always find the most up-to-date version of this document at

<https://www.hepatitisC.uw.edu/go/evaluation-treatment/treatment-goals-predicting-response/core-concept/all>.

Background

The goals for treating persons with chronic hepatitis C virus (HCV) are threefold: (1) eradicate HCV, (2) improve HCV-related health outcomes and survival in all populations, and (3) reduce transmission of HCV to others. For clinicians, the primary and immediate goal is to treat the individual with a regimen that has a very high likelihood of curing the individual of their HCV infection. With the current armamentarium of highly effective and safe direct-acting antiviral (DAA) medications, cure of chronic HCV is expected in more than 95% of persons receiving HCV treatment, regardless of HCV genotype, baseline HCV RNA levels, race, HIV status, or severity of hepatic fibrosis.[1,2] The health outcome benefits following successful treatment of persons with chronic HCV infection are multiple and include reduced prevalence of hepatic fibrosis, lower risk of developing hepatic failure, decreased occurrence of hepatocellular carcinoma (HCC), improved survival, and amelioration of some extrahepatic HCV-related manifestations.[3,4,5,6] With widespread treatment of HCV, the number of persons capable of transmitting HCV would decline dramatically, which could have a major impact on HCV incidence and the overall HCV epidemic.

Virologic Cure and Sustained Virologic Response

HCV Eradication and Sustained Virologic Response (SVR)

The gold standard for determining cure of HCV is the demonstration of sustained undetectable HCV RNA levels after treatment.[\[7\]](#) A sustained virologic response (SVR) is an undetectable HCV RNA level using a sensitive assay (typically with a lower limit of 25 IU/mL) at least 12 weeks after completing HCV therapy ([Figure 1](#)).[\[8\]](#) In the current era, most expert guidelines recommend measuring an HCV RNA level 12 weeks after therapy to evaluate for SVR; individuals with an undetectable HCV RNA level at 12-week posttreatment, also known as an SVR12, are considered to have achieved a virologic cure.[\[7\]](#) Among persons who achieve an SVR12 with direct-acting antiviral (DAA) HCV therapy, greater than 99% go on to achieve a durable response and ongoing absence of detectable viremia.[\[9,10\]](#) Trial outcomes with SVR time frames of shorter duration (SVR8 and SVR4) have been presented in an effort to expedite conclusions from these trials, but SVR4 and SVR8 are not the standard time frame used in clinical practice to determine cure.

Durability of SVR

Long-term follow-up of persons who achieve an SVR12 or SVR24 has shown that nearly 100% remain HCV RNA negative years after therapy.[\[11,12,13\]](#) Several large studies have shown a minimal relapse rate, between 0 to 1% at 5 years.[\[14,15\]](#) Thus, an undetectable HCV RNA 12 or 24 weeks after antiviral therapy can be considered a virologic cure. It is important to note that persons cured of HCV can become reinfected with HCV.[\[15,16\]](#)

Impact of HCV Treatment on Clinical Outcomes

Impact of HCV Treatment on Hepatic Fibrosis

Individuals who achieve an SVR are more likely to have an improvement in liver inflammation and fibrosis than those who do not achieve an SVR.[17,18,19,20] The following studies highlight data related to the impact of HCV eradication on hepatic fibrosis.

- In a pooled analysis of adults who had paired liver biopsies before and 1 month to 6 years after treatment with interferon-based therapies, individuals who achieved an SVR were twice as likely to have lower necroinflammatory scores after treatment, compared to those with virologic relapse (67% versus 32%) and some patients with an SVR had complete regression of liver fibrosis.[20]
- In a meta-analysis, investigators evaluated the impact of HCV treatment on liver stiffness, as measured by vibration-controlled transient elastography.[17] Individuals who achieved an SVR12 had a significantly greater decrease in liver stiffness at the end of treatment and after treatment than patients who failed to achieve an SVR12 (Figure 2).[17] In addition, the decline in liver stiffness among those who achieved an SVR12 was greater with DAA treatment than with interferon-based therapy (decrease of 5.1 kPa versus decrease of 2.8 kPa).[17]
- In a recent study, investigators performed liver stiffness measurements in 70 patients treated with DAA therapy, among whom 95.7% achieved an SVR.[18] Treatment of HCV with DAA therapy resulted in a significant decrease in liver stiffness at the end of treatment and at 12 months posttreatment when compared with baseline measurements (Figure 3).[18]
- In a multi-center, prospective cohort study that enrolled 71 Danish adults with advanced fibrosis, investigators found that liver stiffness decreased by an average of 20% at the end of sofosbuvir-based direct-acting antiviral therapy, and by an additional 15% 1 year after treatment, a finding that was suggestive of fibrosis regression.[21]

Impact of HCV Treatment on Hepatocellular Carcinoma

Considering that achievement of SVR reduces liver fibrosis, one might expect that successful treatment of HCV would lead to a reduced risk of HCC. The following provides a summary of the impact of HCV DAA-based therapy on HCC occurrence and reoccurrence.

- Multiple large, controlled studies have clearly shown a reduction in the risk of HCC occurrence after achievement of SVR with HCV therapy; in these studies, control groups consisted of persons with HCV who were treated but did not achieve an SVR.[4,22,23,24] Several of the more recent studies exclusively involved persons treated with DAA therapy.[23,25,26]
- Although HCV eradication with HCV therapy reduces the risk of HCC occurrence, the risk of developing HCC remains substantial for persons who, prior to treatment, had advanced fibrosis or cirrhosis.[26] Accordingly, individuals who met HCC surveillance criteria prior to HCV treatment should continue to receive HCC surveillance every 6 months after achieving an SVR with HCV treatment.[8,26,27]
- It is important to note posttreatment transient elastography often overestimates the regression in fibrosis, probably because of reduced hepatic inflammation and congestion. Thus, persons with F3-4 fibrosis prior to HCV treatment should continue to have regular hepatocellular carcinoma surveillance, even if testing shows a reduction in liver stiffness or improvement in other non-invasive fibrosis measures.
- Among persons with a history of successfully treated HCC, there are conflicting data regarding the impact of DAA therapy on the risk of HCC recurrence; most studies report no difference in the risk of recurrent HCC among DAA-treated and untreated persons. As such, DAA-treated individuals with a history of HCC require close ongoing HCC surveillance.[27,28,29,30,31]

Impact of HCV Treatment on Survival

In persons with chronic HCV infection, treatment with achievement of SVR12 or SVR24 has been shown to markedly reduce the risk of death, including liver-related and non-liver-related deaths.[[6,14,32,33](#)] Recently, multiple studies have shown major survival benefit in persons with chronic HCV who achieve SVR with DAA therapy.[[34,35](#)] The following summarizes key data related to the impact of DAA therapy on survival after achieving an SVR.

- In a retrospective cohort study, investigators examined the impact of HCV treatment during the years 1990 and 2003 in 5 hepatology units in Europe and Canada.[[36](#)] Individuals with advanced fibrosis who underwent antiviral therapy and achieved an SVR had reduced overall mortality, liver-related death, liver failure, and hepatocellular carcinoma compared with those who did not achieve an SVR ([Figure 4](#)).[[36](#)]
- In a meta-analysis of 35 studies that included 33,360 persons with chronic HCV infection, investigators showed a clear benefit in 5-year overall survival in persons who achieved an SVR with treatment when compared with those who had not achieved an SVR; all the studies analyzed all involved interferon-based therapies, and some included individuals with cirrhosis or persons with HIV coinfection ([Figure 5](#)).[[37](#)]
- In an observational cohort analysis of 103,346 persons with chronic HCV (genotype 1, 2, or 3) in the Veterans Affairs Hepatitis C Clinical Case Registry, investigators examined the impact of achieving SVR with DAA treatment on mortality.[[34](#)] Among the 40,664 persons treated with a DAA regimen, 39,374 (96.8%) achieved an SVR. The mortality rate in persons who achieved an SVR was significantly lower than in those who did not achieve an SVR with treatment, after adjusting for baseline demographics, clinical characteristics, and comorbidity ([Figure 6](#)).[[34](#)] The reduction in mortality was 69.3% among those who achieved an SVR compared to persons who did not receive HCV treatment.[[34](#)]
- Recent cohort studies of persons with HCV and successfully treated HCC have found a significantly lower overall risk of death among persons who receive DAA therapy when compared to those who do not, suggesting that patients with a history of HCC should receive HCV DAA treatment due to multiple treatment-related benefits, including overall improved survival.[[38,39](#)]
- A few studies evaluating contemporary trends have suggested an early signal of benefit since the introduction of DAA therapy with decreased mortality and liver transplantation rates in patients with advanced HCV liver disease.[[40,41](#)]

Impact of HCV Treatment on Extrahepatic Manifestations

Infection with HCV can cause a myriad of extrahepatic complications, including cryoglobulinemia, membranoproliferative glomerulonephritis, dermatologic disorders, insulin resistance and diabetes mellitus, and B-cell non-Hodgkin's lymphoma.[[42,43,44](#)] There is growing evidence that most HCV-related extrahepatic manifestations improve after eradication of HCV, particularly in females.[[45,46](#)] Most notably, a retrospective cohort study that involved 160,875 United States veterans with chronic HCV, patients who achieved an SVR12 with interferon-based therapy had substantial reductions in HCV-related extrahepatic manifestations when compared with individuals who did not achieve an SVR with HCV treatment or were not treated at all ([Figure 7](#)).[[47](#)] In some patients, successful treatment of hepatitis C is associated with improvement or remission of these underlying conditions.[[48,49](#)] In addition, achieving an SVR has been shown to reduce the chance of impaired fasting glucose and diabetes development by 50%, an effect that is independent of other established risk factors for diabetes, such as age and body mass index. A recent meta-analysis of 48 studies showed that an SVR reduced extrahepatic mortality by 56%, improved response to malignant B-cell lymphoproliferative therapy, and vastly improved the chances of a complete resolution of cryoglobulinemic vasculitis.[[3](#)]

Viral Factors that May Impact Response to Therapy

HCV Genotype

Hepatitis C is classified into 6 major genotypes, numbered 1 through 6. In the prior interferon era of treatment, genotype was the strongest predictor of obtaining an SVR.[[50,51,52](#)] In the current direct-acting antiviral (DAA) era, particularly with the approval of pangenotypic regimens, the role of HCV genotype in predicting treatment response has decreased significantly given the high efficacy of different DAA combinations across all genotypes and the introduction of pangenotypic agents. Overall, with a preferred regimen, the SVR12 rate is greater than 95%, regardless of HCV genotype.[[53,54,55,56,57](#)]

HCV RNA Level

In the registration clinical trials of peginterferon and ribavirin, a baseline HCV RNA level over 2 million IU/mL was associated with a 9% lower chance of cure.[[50](#)] Subsequent studies found that patients with high HCV RNA levels and genotype 1 infection had a 16 to 27% lower chance of achieving an SVR depending on the cutoff used.[[58](#)] In the current DAA era, the baseline HCV RNA usually has little impact on the likelihood of achieving an SVR. One exception to this is when considering whether to use 8 or 12 weeks of ledipasvir-sofosbuvir in treatment-naïve persons. A post hoc analysis from the ION-3 trial in treatment-naïve patients without cirrhosis noted that participants with a baseline HCV RNA level less than 6 million IU/mL had similar relapse rates using 8 or 12 weeks of therapy.[[59](#)] Subsequent studies have also shown comparable SVR rates for treatment-naïve individuals without cirrhosis who received either 8 or 12 weeks of ledipasvir-sofosbuvir (if the baseline HCV RNA level was less than 6 million IU/mL).[[60](#)] Baseline HCV RNA levels have not been associated with likelihood of SVR with the more recently approved pangenotypic DAA regimens.

Host Factors that May Impact Response to Therapy

Race

Several clinical trials involving DAA therapy did not demonstrate significant differences in SVR by race, although these trials may not have been sufficiently powered to detect a difference if one existed.[[61,62,63](#)] Several observational studies of DAA effectiveness in the United States Veterans Administration have suggested a slightly reduced likelihood of SVR among Black individuals after adjusting for baseline characteristics.[[64,65,66](#)] The largest of these observational studies analyzed 2014-2015 DAA HCV treatment data by race and found similar overall SVR rates ([Figure 8](#)), but after adjusting for baseline characteristics, the odds of achieving an SVR were slightly lower in Black or Hispanic individuals, when compared with White persons ([Figure 9](#)).[[66](#)] It is important to note that the absolute difference in SVR rates were small (1 to 4 percentage points) and these findings have not been reproduced in settings outside the Veterans Administration. One analysis using nationwide observational data from the Veterans Administration facilities found that among persons with chronic HCV, Black individuals were less likely to receive DAA therapy than White persons.[[67](#)] Overall, when taken together, these data show that SVR12 rates with DAA therapy are excellent across all racial groups, and that while subtle differences might exist, they do not in any way preclude the use of these DAA in Black or Hispanic persons.[[68](#)]

Age

In multiple studies with interferon-based therapy, increasing age was significantly associated with poor tolerance and a lower chance of cure.[[69,70](#)] In contrast, HCV treatment with DAA therapy in the elderly is well tolerated and SVR rates are similar to those seen in younger patients.[[69,71,72](#)] The excellent response in elderly patients has included data in septuagenarians and octogenarians.[[73](#)] In a retrospective analysis of 17,487 patients with chronic HCV infection in the Department of Veterans Affairs healthcare system, age did not significantly impact the SVR rates, even when analyzed using multivariate models ([Figure 10](#)).[[72](#)]

Degree of Hepatic Fibrosis

Advanced fibrosis is typically defined as F3 (pre-cirrhosis or bridging fibrosis) and F4 (cirrhosis) on liver biopsy. In earlier DAA trials, lower SVR rates were observed among persons with compensated cirrhosis.[[64,74](#)] In subsequent trials, newer medication, longer duration of treatment, and modified therapy (with the addition of ribavirin) have all contributed to improved responses in patients with compensated cirrhosis.[[63,75,76,77](#)] The one exception to this has been treatment of persons with genotype 3 HCV and cirrhosis, a group that has emerged as the hardest to treat in the DAA era. Nevertheless, two regimens—glecaprevir-pibrentasvir and sofosbuvir-velpatasvir have been shown to achieve high SVR rates in persons with genotype 3 HCV and compensated cirrhosis.[[56,57](#)] Similarly, when using currently recommended DAA regimens for persons with compensated cirrhosis, studies show SVR12 rates are greater than 90% across all genotypes.[[78](#)] Individuals with decompensated cirrhosis (Child-Turcotte-Pugh class B or C) treated with 12 weeks of ledipasvir-sofosbuvir have lower SVR rates (86 to 87%) compared with SVR rates of 95% or greater in similarly treated persons without cirrhosis.[[79](#)] In a similar study, SVR12 rates of 94% were observed in persons with decompensated cirrhosis when treated with a 12-week regimen of sofosbuvir-velpatasvir plus ribavirin.[[80](#)]

Prior Treatment

Type of Treatment Response with Prior Failure

Prior treatment failure with interferon-based therapy does not significantly impact treatment response to DAA therapy. In general, when using DAA therapy for persons previously treated with an interferon-based regimen, more than 95% should achieve an SVR with retreatment. In contrast, prior treatment failure with a regimen that consisted of or included DAA therapy can impact retreatment response rates, especially if certain HCV resistance-associated substitutions developed during treatment.[\[81,82\]](#) Nevertheless, with the multiple DAA options now available, most persons with prior DAA treatment failure have very good options available.[\[83\]](#) The addition of ribavirin or extension of therapy duration may be required with some regimens to overcome the treatment refractory aspect of persons who previously experienced treatment failure.[\[83,84\]](#)

Summary Points

- There are multiple goals with antiviral therapy for hepatitis C, including (1) eradicate HCV, (2) improve HCV-related health outcomes and survival in all populations, and (3) reduce transmission of HCV to others.
- A sustained virologic response is defined as an undetectable HCV RNA level 12 weeks after stopping antivirals; this is referred to as the SVR12, and the SVR12 has a high correlation with SVR24. An SVR is durable and indicates HCV cure.
- Achieving an SVR following HCV treatment results in improvement of hepatic fibrosis, decreased development of HCC, improvement in survival, and reduction in extrahepatic manifestations associated with chronic HCV.
- In the DAA treatment era, HCV genotype has a reduced role in predicting treatment response given the availability of a variety of DAA combinations with high efficacy across genotypes.
- Older patients, including those 70 years of age and older have comparable responses to DAA therapy when compared with younger patients.
- With newer DAA therapies, individuals with more advanced fibrosis and compensated cirrhosis typically have HCV treatment SVR rates greater than 95% with 12-week treatment regimens. Persons with decompensated cirrhosis are more difficult to treat and often have reduced response rates.

Citations

1. Asselah T, Marcellin P, Schinazi RF. Treatment of hepatitis C virus infection with direct-acting antiviral agents: 100% cure? Liver Int. 2018;38 Suppl 1:7-13.
[\[PubMed Abstract\]](#) -
2. Marcellin F, Roux P, Protopopescu C, Duracinsky M, Spire B, Carrieri MP. Patient-reported outcomes with direct-acting antivirals for the treatment of chronic hepatitis C: current knowledge and outstanding issues. Expert Rev Gastroenterol Hepatol. 2017;11:259-268.
[\[PubMed Abstract\]](#) -
3. Cacoub P, Desbois AC, Comarmond C, Saadoun D. Impact of sustained virological response on the extrahepatic manifestations of chronic hepatitis C: a meta-analysis. Gut. 2018;67:2025-34.
[\[PubMed Abstract\]](#) -
4. Morgan RL, Baack B, Smith BD, Yartel A, Pitasi M, Falck-Ytter Y. Eradication of hepatitis C virus infection and the development of hepatocellular carcinoma: a meta-analysis of observational studies. Ann Intern Med. 2013;158:329-37.
[\[PubMed Abstract\]](#) -
5. Tada T, Kumada T, Toyoda H, et al. Viral eradication reduces all-cause mortality, including non-liver-related disease, in patients with progressive hepatitis C virus-related fibrosis. J Gastroenterol Hepatol. 2017;32:687-694.
[\[PubMed Abstract\]](#) -
6. van der Meer AJ, Veldt BJ, Feld JJ, et al. Association between sustained virological response and all-cause mortality among patients with chronic hepatitis C and advanced hepatic fibrosis. JAMA. 2012;308:2584-93.
[\[PubMed Abstract\]](#) -
7. AASLD-IDSA. HCV Guidance: Recommendations for testing, management, and treating hepatitis C. When and in whom to initiate HCV therapy.
[\[AASLD-IDSA Hepatitis C Guidance\]](#) -
8. AASLD-IDSA. HCV Guidance: Recommendations for testing, management, and treating hepatitis C. Monitoring patients who are starting HCV treatment, are on treatment, or have completed therapy.
[\[AASLD-IDSA HCV Guidance\]](#) -
9. Burgess SV, Hussaini T, Yoshida EM. Concordance of sustained virologic response at weeks 4, 12 and 24 post-treatment of hepatitis c in the era of new oral direct-acting antivirals: A concise review. Ann Hepatol. 2016;15:154-9.
[\[PubMed Abstract\]](#) -
10. Martinot-Peignoux M, Stern C, Maylin S, et al. Twelve weeks posttreatment follow-up is as relevant as 24 weeks to determine the sustained virologic response in patients with hepatitis C virus receiving pegylated interferon and ribavirin. Hepatology. 2010;51:1122-6.
[\[PubMed Abstract\]](#) -
11. George SL, Bacon BR, Brunt EM, Mihindukulasuriya KL, Hoffmann J, Di Bisceglie AM. Clinical, virologic, histologic, and biochemical outcomes after successful HCV therapy: a 5-year follow-up of 150 patients. Hepatology. 2009;49:729-38.
[\[PubMed Abstract\]](#) -

12. Maylin S, Martinot-Peignoux M, Moucari R, et al. Eradication of hepatitis C virus in patients successfully treated for chronic hepatitis C. *Gastroenterology*. 2008;135:821-9.
[[PubMed Abstract](#)] -
13. Morisco F, Granata R, Stroffolini T, et al. Sustained virological response: a milestone in the treatment of chronic hepatitis C. *World J Gastroenterol*. 2013;19:2793-8.
[[PubMed Abstract](#)] -
14. Pearlman BL, Traub N. Sustained virologic response to antiviral therapy for chronic hepatitis C virus infection: a cure and so much more. *Clin Infect Dis*. 2011;52:889-900.
[[PubMed Abstract](#)] -
15. Simmons B, Saleem J, Hill A, Riley RD, Cooke GS. Risk of Late Relapse or Reinfection With Hepatitis C Virus After Achieving a Sustained Virological Response: A Systematic Review and Meta-analysis. *Clin Infect Dis*. 2016;62:683-694.
[[PubMed Abstract](#)] -
16. Page K, Hahn JA, Evans J, et al. Acute hepatitis C virus infection in young adult injection drug users: a prospective study of incident infection, resolution, and reinfection. *J Infect Dis*. 2009;200:1216-26.
[[PubMed Abstract](#)] -
17. Singh S, Facciorusso A, Loomba R, Falck-Ytter YT. Magnitude and Kinetics of Decrease in Liver Stiffness After Antiviral Therapy in Patients With Chronic Hepatitis C: A Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol*. 2018;16:27-38.e4.
[[PubMed Abstract](#)] -
18. Chan J, Gogela N, Zheng H, et al. Direct-Acting Antiviral Therapy for Chronic HCV Infection Results in Liver Stiffness Regression Over 12 Months Post-treatment. *Dig Dis Sci*. 2018;63:486-492.
[[PubMed Abstract](#)] -
19. Poynard T, McHutchison J, Manns M, et al. Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C. *Gastroenterology*. 2002;122:1303-13.
[[PubMed Abstract](#)] -
20. Pockros PJ, Hamzeh FM, Martin P, et al. Histologic outcomes in hepatitis C-infected patients with varying degrees of virologic response to interferon-based treatments. *Hepatology*. 2010;52:1193-200.
[[PubMed Abstract](#)] -
21. Laursen TL, Siggaard CB, Kazankov K, et al. Time-dependent improvement of liver inflammation, fibrosis and metabolic liver function after successful direct-acting antiviral therapy of chronic hepatitis C. *J Viral Hepat*. 2020;27:28-35.
[[PubMed Abstract](#)] -
22. Ioannou GN, Green PK, Berry K. HCV eradication induced by direct-acting antiviral agents reduces the risk of hepatocellular carcinoma. *J Hepatol*. 2017. pii: S0168-8278(17)32273-0.
[[PubMed Abstract](#)] -
23. Calvaruso V, Cabibbo G, Cacciola I, et al. Incidence of Hepatocellular Carcinoma in Patients With HCV-Associated Cirrhosis Treated With Direct-Acting Antiviral Agents. *Gastroenterology*. 2018;155:411-21.e4.
[[PubMed Abstract](#)] -
24. Nahon P, Bourcier V, Layese R, et al. Eradication of Hepatitis C Virus Infection in Patients With Cirrhosis Reduces Risk of Liver and Non-Liver Complications. *Gastroenterology*. 2017;152:142-156.e2.

[\[PubMed Abstract\]](#) -

25. Konjeti VR, John BV. Interaction Between Hepatocellular Carcinoma and Hepatitis C Eradication With Direct-acting Antiviral Therapy. *Curr Treat Options Gastroenterol*. 2018;16:203-214.
[\[PubMed Abstract\]](#) -
26. Kanwal F, Kramer J, Asch SM, Chayanupatkul M, Cao Y, El-Serag HB. Risk of Hepatocellular Cancer in HCV Patients Treated With Direct-Acting Antiviral Agents. *Gastroenterology*. 2017;153:996-1005.e1.
[\[PubMed Abstract\]](#) -
27. Singal AG, Lim JK, Kanwal F. AGA Clinical Practice Update on Interaction Between Oral Direct-Acting Antivirals for Chronic Hepatitis C Infection and Hepatocellular Carcinoma: Expert Review. *Gastroenterology*. 2019;156:2149-57.
[\[PubMed Abstract\]](#) -
28. ANRS collaborative study group on hepatocellular carcinoma (ANRS CO22 HEPATHER, CO12 CirVir and CO23 CUPILT cohorts). . Lack of evidence of an effect of direct-acting antivirals on the recurrence of hepatocellular carcinoma: Data from three ANRS cohorts. *J Hepatol*. 2016;65:734-40.
[\[PubMed Abstract\]](#) -
29. Cabibbo G, Petta S, Calvaruso V, et al. Is early recurrence of hepatocellular carcinoma in HCV cirrhotic patients affected by treatment with direct-acting antivirals? A prospective multicentre study. *Aliment Pharmacol Ther*. 2017;46:688-695.
[\[PubMed Abstract\]](#) -
30. Saraiya N, Yopp AC, Rich NE, Odewole M, Parikh ND, Singal AG. Systematic review with meta-analysis: recurrence of hepatocellular carcinoma following direct-acting antiviral therapy. *Aliment Pharmacol Ther*. 2018;48:127-37.
[\[PubMed Abstract\]](#) -
31. Singal AG, Rich NE, Mehta N, et al. Direct-Acting Antiviral Therapy Not Associated With Recurrence of Hepatocellular Carcinoma in a Multicenter North American Cohort Study. *Gastroenterology*. 2019;156:1683-92.e1.
[\[PubMed Abstract\]](#) -
32. Backus LI, Boothroyd DB, Phillips BR, Belperio P, Halloran J, Mole LA. A sustained virologic response reduces risk of all-cause mortality in patients with hepatitis C. *Clin Gastroenterol Hepatol*. 2011;9:509-516.e1.
[\[PubMed Abstract\]](#) -
33. Berenguer J, Rodríguez E, Miralles P, et al. Sustained virological response to interferon plus ribavirin reduces non-liver-related mortality in patients coinfectd with HIV and Hepatitis C virus. *Clin Infect Dis*. 2012;55:728-36.
[\[PubMed Abstract\]](#) -
34. Backus LI, Belperio PS, Shahoumian TA, Mole LA. Direct-acting antiviral sustained virologic response: Impact on mortality in patients without advanced liver disease. *Hepatology*. 2018;68:827-38.
[\[PubMed Abstract\]](#) -
35. Backus LI, Belperio PS, Shahoumian TA, Mole LA. Impact of Sustained Virologic Response with Direct-Acting Antiviral Treatment on Mortality in Patients with Advanced Liver Disease. *Hepatology*. 2019;69:487-97.
[\[PubMed Abstract\]](#) -

36. Veldt BJ, Heathcote EJ, Wedemeyer H, et al. Sustained virologic response and clinical outcomes in patients with chronic hepatitis C and advanced fibrosis. *Ann Intern Med.* 2007;147:677-84.
[\[PubMed Abstract\]](#) -
37. Simmons B, Saleem J, Heath K, Cooke GS, Hill A. Long-Term Treatment Outcomes of Patients Infected With Hepatitis C Virus: A Systematic Review and Meta-analysis of the Survival Benefit of Achieving a Sustained Virological Response. *Clin Infect Dis.* 2015;61:730-40.
[\[PubMed Abstract\]](#) -
38. Cabibbo G, Celsa C, Calvaruso V, et al. Direct-acting antivirals after successful treatment of early hepatocellular carcinoma improve survival in HCV-cirrhotic patients. *J Hepatol.* 2019;71:265-73.
[\[PubMed Abstract\]](#) -
39. Singal AG, Rich NE, Mehta N, et al. Direct-Acting Antiviral Therapy for Hepatitis C Virus Infection Is Associated With Increased Survival in Patients With a History of Hepatocellular Carcinoma. *Gastroenterology.* 2019;157:1253-63.e2.
[\[PubMed Abstract\]](#) -
40. Arora SS, Axley P, Ahmed Z, et al. Decreasing frequency and improved outcomes of hepatitis C-related liver transplantation in the era of direct-acting antivirals - a retrospective cohort study. *Transpl Int.* 2019;32:854-64.
[\[PubMed Abstract\]](#) -
41. Kim D, Li AA, Perumpail BJ, et al. Changing Trends in Etiology-Based and Ethnicity-Based Annual Mortality Rates of Cirrhosis and Hepatocellular Carcinoma in the United States. *Hepatology.* 2019;69:1064-74.
[\[PubMed Abstract\]](#) -
42. El-Serag HB, Hampel H, Yeh C, Rabeneck L. Extrahepatic manifestations of hepatitis C among United States male veterans. *Hepatology.* 2002;36:1439-45.
[\[PubMed Abstract\]](#) -
43. Jacobson IM, Cacoub P, Dal Maso L, Harrison SA, Younossi ZM. Manifestations of chronic hepatitis C virus infection beyond the liver. *Clin Gastroenterol Hepatol.* 2010;8:1017-29.
[\[PubMed Abstract\]](#) -
44. Tang L, Marcell L, Kottlilil S. Systemic manifestations of hepatitis C infection. *Infect Agent Cancer.* 2016;11:29.
[\[PubMed Abstract\]](#) -
45. Ioannou GN, Feld JJ. What Are the Benefits of a Sustained Virologic Response to Direct-Acting Antiviral Therapy for Hepatitis C Virus Infection? *Gastroenterology.* 2019;156:446-460.e2.
[\[PubMed Abstract\]](#) -
46. Li J, Gordon SC, Zhou Y, et al. Sex Differences in Extrahepatic Outcomes After Antiviral Treatment for Hepatitis C. *Am J Gastroenterol.* 2021;116:576-83.
[\[PubMed Abstract\]](#) -
47. Mahale P, Engels EA, Li R, et al. The effect of sustained virological response on the risk of extrahepatic manifestations of hepatitis C virus infection. *Gut.* 2018;67:553-561.
[\[PubMed Abstract\]](#) -
48. Arase Y, Suzuki F, Suzuki Y, et al. Sustained virological response reduces incidence of onset of type 2 diabetes in chronic hepatitis C. *Hepatology.* 2009;49:739-44.

[\[PubMed Abstract\]](#) -

49. Conjeevaram HS, Wahed AS, Afdhal N, Howell CD, Everhart JE, Hoofnagle JH. Changes in insulin sensitivity and body weight during and after peginterferon and ribavirin therapy for hepatitis C. *Gastroenterology*. 2010;140:469-77.
[\[PubMed Abstract\]](#) -
50. Fried MW, Shiffman ML, Reddy KR, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med*. 2002;347:975-82.
[\[PubMed Abstract\]](#) -
51. Hadziyannis SJ, Sette H Jr, Morgan TR, et al. Peginterferon-alfa2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med*. 2004;140:346-55.
[\[PubMed Abstract\]](#) -
52. Manns MP, McHutchison JG, Gordon SC, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet*. 2001;358:958-65.
[\[PubMed Abstract\]](#) -
53. Zeuzem S, Foster GR, Wang S, et al. Glecaprevir-Pibrentasvir for 8 or 12 Weeks in HCV Genotype 1 or 3 Infection. *N Engl J Med*. 2018;378:354-369.
[\[PubMed Abstract\]](#) -
54. Feld JJ, Foster GR. Second generation direct-acting antivirals - Do we expect major improvements? *J Hepatol*. 2016;65:S130-S142.
[\[PubMed Abstract\]](#) -
55. Falade-Nwulia O, Suarez-Cuervo C, Nelson DR, Fried MW, Segal JB, Sulkowski MS. Oral Direct-Acting Agent Therapy for Hepatitis C Virus Infection: A Systematic Review. *Ann Intern Med*. 2017;166:637-48.
[\[PubMed Abstract\]](#) -
56. Foster GR, Afdhal N, Roberts SK, et al. Sofosbuvir and velpatasvir for HCV genotype 2 and 3 infection. *N Engl J Med*. 2015;373:2608-17.
[\[PubMed Abstract\]](#) -
57. Wyles D, Poordad F, Wang S, et al. Glecaprevir/pibrentasvir for hepatitis C virus genotype 3 patients with cirrhosis and/or prior treatment experience: A partially randomized phase 3 clinical trial. *Hepatology*. 2018;67:514-23.
[\[PubMed Abstract\]](#) -
58. Zeuzem S, Rodríguez-Torres M, Rajender Reddy K, et al. Optimized threshold for serum HCV RNA to predict treatment outcomes in hepatitis C patients receiving peginterferon alfa-2a/ribavirin. *J Viral Hepat*. 2012;19:766-74.
[\[PubMed Abstract\]](#) -
59. Kowdley KV, Gordon SC, Reddy KR, et al. Ledipasvir and sofosbuvir for 8 or 12 weeks for chronic HCV without cirrhosis. *N Engl J Med*. 2014;370:1879-88.
[\[PubMed Abstract\]](#) -
60. Terrault NA, Zeuzem S, Di Bisceglie AM, et al. Effectiveness of Ledipasvir-Sofosbuvir Combination in Patients With Hepatitis C Virus Infection and Factors Associated With Sustained Virologic Response. *Gastroenterology*. 2016;151:1131-1140.e5.

[\[PubMed Abstract\]](#) -

61. Roth D, Nelson DR, Bruchfeld A, et al. Grazoprevir plus elbasvir in treatment-naïve and treatment-experienced patients with hepatitis C virus genotype 1 infection and stage 4-5 chronic kidney disease (the C-SURFER study): a combination phase 3 study. *Lancet*. 2015;386:1537-45.
[\[PubMed Abstract\]](#) -
62. Wilder JM, Jeffers LJ, Ravendhran N, et al. Safety and efficacy of ledipasvir-sofosbuvir in black patients with hepatitis C virus infection: A retrospective analysis of phase 3 data. *Hepatology*. 2016;63:437-44.
[\[PubMed Abstract\]](#) -
63. Zeuzem S, Ghalib R, Reddy KR, et al. Grazoprevir-Elbasvir Combination Therapy for Treatment-Naïve Cirrhotic and Noncirrhotic Patients With Chronic Hepatitis C Virus Genotype 1, 4, or 6 Infection: A Randomized Trial. *Ann Intern Med*. 2015;163:1-13.
[\[PubMed Abstract\]](#) -
64. Ioannou GN, Beste LA, Chang MF, et al. Effectiveness of Sofosbuvir, Ledipasvir/Sofosbuvir, or Paritaprevir/Ritonavir/Ombitasvir and Dasabuvir Regimens for Treatment of Patients With Hepatitis C in the Veterans Affairs National Health Care System. *Gastroenterology*. 2016;151:457-471.e5.
[\[PubMed Abstract\]](#) -
65. Benhammou JN, Dong TS, May FP, et al. Race affects SVR12 in a large and ethnically diverse hepatitis C-infected patient population following treatment with direct-acting antivirals: Analysis of a single-center Department of Veterans Affairs cohort. *Pharmacol Res Perspect*. 2018;6:e00379.
[\[PubMed Abstract\]](#) -
66. Su F, Green PK, Berry K, Ioannou GN. The association between race/ethnicity and the effectiveness of direct antiviral agents for hepatitis C virus infection. *Hepatology*. 2017;65:426-438.
[\[PubMed Abstract\]](#) -
67. Kanwal F, Kramer JR, El-Serag HB, et al. Race and Gender Differences in the Use of Direct Acting Antiviral Agents for Hepatitis C Virus. *Clin Infect Dis*. 2016;63:291-9.
[\[PubMed Abstract\]](#) -
68. Wilder JM, Muir A. Direct acting antivirals for the treatment of hepatitis C in ethnic minority populations. *Expert Opin Pharmacother*. 2018;19:451-456.
[\[PubMed Abstract\]](#) -
69. Rheem J, Sundaram V, Saab S. Antiviral Therapy in Elderly Patients With Hepatitis C Virus Infection. *Gastroenterol Hepatol (N Y)*. 2015;11:294-346.
[\[PubMed Abstract\]](#) -
70. Huang CF, Yang JF, Dai CY, et al. Efficacy and safety of pegylated interferon combined with ribavirin for the treatment of older patients with chronic hepatitis C. *J Infect Dis*. 2010;201:751-9.
[\[PubMed Abstract\]](#) -
71. Huang CF, Yu ML. Treating hepatitis C in the elderly: pharmacotherapeutic considerations and developments. *Expert Opin Pharmacother*. 2017;18:1867-1874.
[\[PubMed Abstract\]](#) -
72. Su F, Beste LA, Green PK, Berry K, Ioannou GN. Direct-acting antivirals are effective for chronic hepatitis C treatment in elderly patients: a real-world study of 17 487 patients. *Eur J Gastroenterol Hepatol*. 2017;29:686-693.
[\[PubMed Abstract\]](#) -

73. Snyder HS, Ali B, Gonzalez HC, Nair S, Satapathy SK. Efficacy and Safety of Sofosbuvir-Based Direct Acting Antivirals for Hepatitis C in Septuagenarians and Octogenarians. *J Clin Exp Hepatol*. 2017;7:93-96.
[[PubMed Abstract](#)] -
74. Sterling RK, Kuo A, Rustgi VK, et al. Virological outcomes and treatment algorithms utilisation in observational study of patients with chronic hepatitis C treated with boceprevir or telaprevir. *Aliment Pharmacol Ther*. 2015;41:671-85.
[[PubMed Abstract](#)] -
75. Lawitz E, Gane E, Pearlman B, et al. Efficacy and safety of 12 weeks versus 18 weeks of treatment with grazoprevir (MK-5172) and elbasvir (MK-8742) with or without ribavirin for hepatitis C virus genotype 1 infection in previously untreated patients with cirrhosis and patients with previous null response with or without cirrhosis (C-WORTHY): a randomised, open-label phase 2 trial. *Lancet*. 2015;385:1075-86.
[[PubMed Abstract](#)] -
76. Forns X, Lee SS, Valdes J, et al. Glecaprevir plus pibrentasvir for chronic hepatitis C virus genotype 1, 2, 4, 5, or 6 infection in adults with compensated cirrhosis (EXPEDITION-1): a single-arm, open-label, multicentre phase 3 trial. *Lancet Infect Dis*. 2017;17:1062-1068.
[[PubMed Abstract](#)] -
77. Asselah T, Bourgeois S, Pianko S, et al. Sofosbuvir/velpatasvir in patients with hepatitis C virus genotypes 1-6 and compensated cirrhosis or advanced fibrosis. *Liver Int*. 2018;38:443-450.
[[PubMed Abstract](#)] -
78. Kwo PY. Regimens for Cirrhotic Patients. *Clin Liver Dis*. 2015;19:657-67.
[[PubMed Abstract](#)] -
79. Charlton M, Everson GT, Flamm SL, et al. Ledipasvir and sofosbuvir plus ribavirin for treatment of HCV infection in patients with advanced liver disease. *Gastroenterology*. 2015;149:649-59.
[[PubMed Abstract](#)] -
80. Curry MP, O'Leary JG, Bzowej N, et al. Sofosbuvir and Velpatasvir for HCV in Patients with Decompensated Cirrhosis. *N Engl J Med*. 2015;373:2618-28.
[[PubMed Abstract](#)] -
81. Kwo PY, Badshah MB. Treatment of HCV in Patients who Failed First-Generation PI Therapy: a Review of Current Literature. *Curr Gastroenterol Rep*. 2015;17:462.
[[PubMed Abstract](#)] -
82. Wyles DL. Resistance to DAAs: When to Look and When It Matters. *Curr HIV/AIDS Rep*. 2017;14:229-237.
[[PubMed Abstract](#)] -
83. Zuckerman A, Chastain CA, Naggie S. Retreatment Options Following HCV Direct Acting Antiviral Failure. *Curr Treat Options Infect Dis*. 2017;9:389-402.
[[PubMed Abstract](#)] -
84. AASLD-IDSA. HCV Guidance: Recommendations for testing, management, and treating hepatitis C. Retreatment of persons in whom prior therapy failed.
[[AASLD-IDSA Hepatitis C Guidance](#)] -

References

- Andersen ES, Moessner BK, Christensen PB, et al. Lower liver stiffness in patients with sustained virological response 4 years after treatment for chronic hepatitis C. *Eur J Gastroenterol Hepatol*. 2011;23:41-4.
[\[PubMed Abstract\]](#) -
- Butt AA, Wang X, Moore CG. Effect of hepatitis C virus and its treatment on survival. *Hepatology*. 2009;50:387-92.
[\[PubMed Abstract\]](#) -
- Cabibbo G, Petta S, Calvaruso V, et al. Is early recurrence of hepatocellular carcinoma in HCV cirrhotic patients affected by treatment with direct-acting antivirals? A prospective multicentre study. *Aliment Pharmacol Ther*. 2017;46:688-95.
[\[PubMed Abstract\]](#) -
- Campos-Varela I, Castells L, Esteban JI, et al. Twelve-week posttreatment follow-up to predict sustained virologic response for recurrent hepatitis C infection in liver recipients. *Transplantation*. 2012;93:450-3.
[\[PubMed Abstract\]](#) -
- Chang CY, Nguyen P, Le A, et al. Real-world experience with interferon-free, direct acting antiviral therapies in Asian Americans with chronic hepatitis C and advanced liver disease. *Medicine (Baltimore)*. 2017;96:e6128.
[\[PubMed Abstract\]](#) -
- Cheung MCM, Walker AJ, Hudson BE, et al. Outcomes after successful direct-acting antiviral therapy for patients with chronic hepatitis C and decompensated cirrhosis. *J Hepatol*. 2016;65:741-747.
[\[PubMed Abstract\]](#) -
- Dieperink E, Pocha C, Thuras P, Knott A, Colton S, Ho SB. All-cause mortality and liver-related outcomes following successful antiviral treatment for chronic hepatitis C. *Dig Dis Sci*. 2014;59:872-80.
[\[PubMed Abstract\]](#) -
- El-Sherif O, Jiang ZG, Tapper EB, et al. Baseline Factors Associated With Improvements in Decompensated Cirrhosis After Direct-Acting Antiviral Therapy for Hepatitis C Virus Infection. *Gastroenterology*. 2018;154:2111-2121.e8.
[\[PubMed Abstract\]](#) -
- Fernández-Montero JV, Aspinall EJ, Burns JE. Treatment as Prevention: Should Hepatitis C Learn the Lessons from HIV? *AIDS Rev*. 2017;19:212-218.
[\[PubMed Abstract\]](#) -
- Foster GR, Irving WL, Cheung MC, et al. Impact of direct acting antiviral therapy in patients with chronic hepatitis C and decompensated cirrhosis. *J Hepatol*. 2016;64:1224-31.
[\[PubMed Abstract\]](#) -
- Ge D, Fellay J, Thompson AJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature*. 2009;461:399-401.
[\[PubMed Abstract\]](#) -
- Ghany MG, Strader DB, Thomas DL, Seeff LB; American Association for the Study of Liver Diseases. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology*. 2009;49:1335-74.

[\[AASLD Practice Guidelines\]](#) -

- Kutala BK, Guedj J, Asselah T, et al. Impact of treatment against hepatitis C virus on overall survival of naive patients with advanced liver disease. *Antimicrob Agents Chemother.* 2014;59:803-10.
[\[PubMed Abstract\]](#) -
- Lawitz E, Lalezari JP, Hassanein T, et al. Sofosbuvir in combination with peginterferon alfa-2a and ribavirin for non-cirrhotic, treatment-naïve patients with genotypes 1, 2, and 3 hepatitis C infection: a randomised, double-blind, phase 2 trial. *Lancet Infect Dis.* 2013;13:401-8.
[\[PubMed Abstract\]](#) -
- Li DK, Chung RT. Impact of hepatitis C virus eradication on hepatocellular carcinogenesis. *Cancer.* 2015;121:2874-82.
[\[PubMed Abstract\]](#) -
- Mira JA, Rivero-Juárez A, López-Cortés LF, et al. Benefits from sustained virologic response to pegylated interferon plus ribavirin in HIV/hepatitis C virus-coinfected patients with compensated cirrhosis. *Clin Infect Dis.* 2013;56:1646-53.
[\[PubMed Abstract\]](#) -
- Moon C, Jung KS, Kim do Y, et al. Lower incidence of hepatocellular carcinoma and cirrhosis in hepatitis C patients with sustained virological response by pegylated interferon and ribavirin. *Dig Dis Sci.* 2014;60:573-81.
[\[PubMed Abstract\]](#) -
- Muir AJ, Bornstein JD, Killenberg PG. Peginterferon alfa-2b and ribavirin for the treatment of chronic hepatitis C in blacks and non-Hispanic whites. *N Engl J Med.* 2004;350:2265-71.
[\[PubMed Abstract\]](#) -
- Olafsson S, Tyrfinngsson T, Runarsdottir V, et al. Treatment as Prevention for Hepatitis C (TraP Hep C) - a nationwide elimination programme in Iceland using direct-acting antiviral agents. *J Intern Med.* 2018;283:500-7.
[\[PubMed Abstract\]](#) -
- Pol S, Parlati L. Treatment of hepatitis C: the use of the new pangenotypic direct-acting antivirals in "special populations". *Liver Int.* 2018 Feb;38 Suppl 1:28-33.
[\[PubMed Abstract\]](#) -
- Reig M, Mariño Z, Perelló C, et al. Unexpected high rate of early tumor recurrence in patients with HCV-related HCC undergoing interferon-free therapy. *J Hepatol.* 2016;65:719-726.
[\[PubMed Abstract\]](#) -
- Rodriguez-Torres M, Jeffers LJ, Sheikh MY, et al. Peginterferon alfa-2a and ribavirin in Latino and non-Latino whites with hepatitis C. *N Engl J Med.* 2009;360:257-67.
[\[PubMed Abstract\]](#) -
- Smith-Palmer J, Cerri K, Valentine W. Achieving sustained virologic response in hepatitis C: a systematic review of the clinical, economic and quality of life benefits. *BMC Infect Dis.* 2015;15:19.
[\[PubMed Abstract\]](#) -
- Stättermayer AF, Stauber R, Hofer H, et al. Impact of IL28B genotype on the early and sustained virologic response in treatment-naïve patients with chronic hepatitis C. *Clin Gastroenterol Hepatol.* 2011;9:344-50.
[\[PubMed Abstract\]](#) -

- Toccaceli F, Laghi V, Capurso L, Koch M, Sereno S, Scuderi M, Italian HeparNet Group. Long-term liver histology improvement in patients with chronic hepatitis C and sustained response to interferon. J Viral Hepat. 2003;10:126-33.
[\[PubMed Abstract\]](#) -
- Waziry R, Hajarizadeh B, Grebely J, et al. Hepatocellular carcinoma risk following direct-acting antiviral HCV therapy: A systematic review, meta-analyses, and meta-regression. J Hepatol. 2017;67:1204-1212.
[\[PubMed Abstract\]](#) -
- Zelenev A, Li J, Mazhnaya A, Basu S, Altice FL. Hepatitis C virus treatment as prevention in an extended network of people who inject drugs in the USA: a modelling study. Lancet Infect Dis. 2018;18:215-24.
[\[PubMed Abstract\]](#) -

Figures

Figure 1 Sustained Virologic Response 12 (SVR 12)

This graphic shows an example of an SVR12 in a person who received 12 weeks of HCV treatment. The SVR12 is shown by the undetectable HCV RNA 12 weeks after treatment was stopped.

Source: illustration by David H. Spach, MD




Figure 2 Liver Stiffness in Persons With or Without SVR12

In this review and meta-analysis, investigators compared liver stiffness measured by vibration-controlled transient elastography (VCTE) before and after treatment. This graph compares liver stiffness before and after treatment in adults who attained SVR12 with those who do not achieve SVR12. The decline in liver stiffness among persons who achieved an SVR12 was greater when treated with direct-acting antiviral therapy than with interferon-based therapy (decrease of 5.1 kPa versus decrease of 2.8kPa).

Source: Singh S, Facciorusso A, Loomba R, Falck-Ytter YT. Magnitude and Kinetics of Decrease in Liver Stiffness After Antiviral Therapy in Patients With Chronic Hepatitis C: A Systematic Review and Meta-analysis. Clin Gastroenterol Hepatol. 2018;16:27-38.e4.




Figure 3 Liver Stiffness Treatments in Persons Treated with Direct-Acting Antiviral Therapy

This study enrolled 70 persons who received direct-acting antiviral therapy for chronic HCV infection. This graphic shows liver stiffness measurement at baseline, end-of-treatment, and 12-month posttreatment. Overall, 48.6% of the participants had a 30% or greater improvement in the liver stiffness measurement (at end of follow-up compared with baseline).

Source: Chan J, Gogela N, Zheng H, et al. Direct-Acting Antiviral Therapy for Chronic HCV Infection Results in Liver Stiffness Regression Over 12 Months Post-treatment. Dig Dis Sci. 2018;63:486-92.




Figure 4 Clinical Outcome by Response to Treatment in Patients with Chronic Hepatitis C and Advanced Fibrosis

This retrospective study was performed in Europe and Canada and examined whether sustained virologic response following hepatitis C treatment correlated with clinical outcomes. The major finding was that treatment was associated with improved clinical outcomes, primarily because of lower rates of liver failure.


Source: Veldt BJ, Heathcote EJ, Wedemeyer H, et al. Sustained virologic response and clinical outcomes in patients with chronic hepatitis C and advanced fibrosis. Ann Intern Med. 2007;147:677-84.

Clinical Outcomes by Response to Hepatitis C Treatment			
Outcome	Patients with SVR	Patients without SVR	Hazard Ratio
	Events per 10,000 Patient-Years		
Overall death	71	193	0.44
Liver-Related death	36	283	0.14
Non-liver-related Death	36	40	1.21
Liver failure	0	365	0.03
Hepatocellular carcinoma	107	277	0.46

Figure 5 5-Year Survival Rate following HCV Treatment Based on SVR Response

This graphic is based on data from 31 studies published from 2000 to 2014 that included 33,360 participants. The 5-year mortality rates shown are based on whether the patient achieved an SVR.

Source: Simmons B, Saleem J, Heath K, Cooke GS, Hill A. Long-Term Treatment Outcomes of Patients Infected With Hepatitis C Virus: A Systematic Review and Meta-analysis of the Survival Benefit of Achieving a Sustained Virological Response. Clin Infect Dis. 2015;61:730-40.

5-Year Mortality Rates for SVR versus Non-SVR Groups

Abbreviations: SVR= sustained virologic response

Figure 6 Impact of SVR on Mortality Rates with DAA Therapy

Source: Backus LI, Belperio PS, Shahoumian TA, Mole LA. Direct-Acting Antiviral Sustained Virologic Response: Impact on Mortality in Patients without Advanced Liver Disease. *Hepatology*. 2018 Jan 29. [Epub ahead of print]




Figure 7 HCV Treatment and Outcome of Extrahepatic Manifestations


Source: Mahale P, Engels EA, Li R, et al. The effect of sustained virological response on the risk of extrahepatic manifestations of hepatitis C virus infection. Gut. 2018;67:553-61.

Clinical Outcomes by HCV Treatment and Response			
Outcomes	No Treatment	Treatment without SVR	Treatment with SVR
	Events per 1,000 Patient-Years		
Mixed cryoglobulinemia	0.72	0.52	0.33
Glomerulonephritis	2.83	1.62	1.09
Porphyria cutanea tarda	0.52	0.37	0.16
Lichen planus	0.68	0.71	0.56
Non-Hodgkin's lymphoma	0.91	0.55	0.43
Diabetes mellitus	21.6	17.0	13.9
Coronary heart disease	1.01	0.58	0.75
Stroke	9.14	4.64	5.10

Figure 8 Sustained Virologic Response Rates (Unadjusted), by Race

This graphic shows unadjusted sustained virologic response rates with DAA therapy among persons who are White, Black, Hispanic, and persons of another race.

Source: Su F, Green PK, Berry K, Ioannou GN. The association between race/ethnicity and the effectiveness of direct antiviral agents for hepatitis C virus infection. Hepatology. 2017;65:426-38.




Other = Asian, Pacific Islander, American Indian, Alaskan Native

Figure 9 Sustained Virologic Response Rates (Adjusted Odds Ratio), by Race

This graphic shows adjusted odds ratio sustained virologic response rates with DAA therapy among persons who are White, Black, Hispanic, and persons of another race. This model includes the following baseline characteristics: age, genotype/subgenotype, regimen, gender, HCV viral load, platelet count, serum bilirubin level, serum albumin level, alcohol use disorder, diabetes, cirrhosis, decompensated cirrhosis, HCC, liver transplantation, and prior treatment.

Source: Su F, Green PK, Berry K, Ioannou GN. The association between race/ethnicity and the effectiveness of direct antiviral agents for hepatitis C virus infection. *Hepatology*. 2017;65:426-38.



Other = Asian, Pacific Islander, American Indian, Alaskan Native

Figure 10 Sustained Virologic Response, by Age Group

In this retrospective analysis of DAA treatment of HCV in the Department of Veterans Affairs health care system, investigators analyzed treatment response based on age. As shown in this graph, excellent SVR rates occurred across all age groups, including those 75 years of age and older.

Source: Su F, Beste LA, Green PK, Berry K, Ioannou GN. Direct-acting antivirals are effective for chronic hepatitis C treatment in elderly patients: a real-world study of 17 487 patients. *Eur J Gastroenterol Hepatol*. 2017;29:686-93.

